

A proximal DC approach for quadratic assignment problem

Zhuoxuan Jiang · Xinyuan Zhao · Chao Ding

Abstract In this paper, we show that the quadratic assignment problem (QAP) can be reformulated to an equivalent rank constrained doubly nonnegative (DNN) problem. Under the framework of the difference of convex functions (DC) approach, a semi-proximal DC algorithm (DCA) is proposed for solving the relaxation of the rank constrained DNN problem whose subproblems can be solved by the semi-proximal augmented Lagrangian method (sPALM). We show that the generated sequence converges to a stationary point of the corresponding DC problem, which is feasible to the rank constrained DNN problem under some suitable assumptions. Moreover, numerical experiments demonstrate that for most QAP instances, the proposed approach can find the global optimal solutions efficiently, and for others, the proposed algorithm is able to provide good feasible solutions in a reasonable time.

Keywords quadratic assignment problem · doubly nonnegative programming · augmented Lagrangian method · rank constraint

Mathematics Subject Classification (2010) 90C22 · 90C25 · 90C26 · 90C27

Z. X. Jiang
College of Applied Sciences, Beijing University of Technology, Beijing, P.R. China.
E-mail: zxjiang@emails.bjut.edu.cn

X. Y. Zhao
College of Applied Sciences, Beijing University of Technology, Beijing, P.R. China. The research of this author was supported by the National Natural Science Foundation of China under projects No. 11871002 and the General Program of Science and Technology of Beijing Municipal Education Commission No. KM201810005004.
E-mail: xyzhao@bjut.edu.cn

C. Ding
Institute of Applied Mathematics, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing, P.R. China. The research of this author was supported by the National Natural Science Foundation of China under projects No. 11671387, No. 11531014 and No. 11688101 and the Beijing Natural Science Foundation (Z190002).
E-mail: dingchao@amss.ac.cn

1 Introduction

The quadratic assignment problem (QAP) is a classical mathematical model for location theory, which is used to model the location problem of allocating n facilities to n locations while minimizing the quadratic objective coming from the distance between the locations and the flow between the facilities. The standard form introduced by Koopmans and Beckmann [27] is as following:

$$\min \left\{ \sum_{1 \leq i, j \leq n} A_{ij} B_{\pi(i), \pi(j)} + \sum_i C_{i\pi(i)} \mid \pi \in \mathcal{P}^n \right\}, \quad (1)$$

where A , B and C are given $n \times n$ real matrices and \mathcal{P}^n is the the group of all permutations of $\{1, \dots, n\}$. In this paper, we make the standard assumption that A and B are symmetric.

Nowadays, QAP becomes one of the most important combinatorial optimization problems due to [its wide applications](#) in many different areas, such as chip design, manufacturing, computer graphics and vision, and others (see [14, 17] for more details). However, it is well known that QAP is NP-hard [39] and still quite difficult to compute the problems of dimension $n \geq 30$ in a reasonable computational time. Exact solution algorithms for QAP in practice are usually based on the branch and bound technique used to reduce the domain and improve the bounds of relaxation problems [3]. Therefore, it is still an important research topic to improve the lower or upper bounds for QAP efficiently.

Meanwhile, semidefinite programming (SDP) [44] has proven to be very successful in this trend by providing tight relaxations for hard combinatorial problems [45]. To obtain lower bounds for QAP, various SDP relaxations are established [29, 50]. Although SDP relaxation is numerically successful, it does not satisfy the Slater condition that may make the dual optimal solution unbounded [36]. That is an important reason why some interior-point methods become inefficient for solving QAPs. To overcome this difficulty, by exploring the geometrical structure of SDP relaxations, Zhao et al. [50] considered a reduced SDP problem by projecting the primal problem onto the minimal face of the semidefinite cone, and constructed some Slater points for such SDP relaxations, which can be solved by the interior-point method and the bundle method [37] efficiently for $n \leq 30$.

To improve the quality of the SDP relaxation of QAP, Povh and Rendl [35] showed that the optimal value of QAP was equal to the optimal value of the convex completely positive programming (CPP), i.e., a linear program over the cone of completely positive matrices. In fact, based on [13], many important binary and nonconvex quadratic programs including QAP can be equivalently reformulated as the convex CPPs, under some mild conditions. However, these CPP reformulations are known to be numerically intractable [33], and an efficient strategy is replacing the completely positive cone with doubly nonnegative (DNN) cone and solving the relaxation problems by SDP solvers [20, 26, 46, 48, 49, 51]. The QAP and the corresponding CPP relaxation

proposed by Povh and Rendl [35] have the same optimal value, but the optimal solution may be different except that the rank of the optimal solution is one. Because it is well-known that the rank constrained matrix optimization problems are computationally intractable and difficult in general [15], the rank one constraints are usually dropped in both the CPP and its related DNN relaxations of QAP. However, by using the difference of two convex functions (DC), the rank constraint can be replaced by the difference of the nuclear norm function and Ky-Fan k -norm function. Based on this simple observation, a penalty approach is proposed by [21] for calibrating rank constrained correlation matrix problems, which usually performs very well in many applications (see also [28]). Based on the DC reformulations of the rank constraints, we shall reformulate the original QAP as a DC programming [1, 2] and employ the DC algorithm (DCA) to solve the nonconvex QAP relaxation problems.

In this paper, we will propose a new rank constrained DNN model and show that it is equivalent to the original QAP (in the sense of both optimal values and optimal solutions). Also, we shall show the same techniques can be applied to other important nonconvex problems such as the standard quadratic programming and the minimum-cut graph tri-partitioning problem. Although the equivalent rank constrained DNN model is still numerically intractable, we will propose a semi-proximal DC algorithm (DCA) framework for finding a feasible stationary point. Furthermore, for the large-scaled DCA inner subproblems, we will apply an efficient majorized semismooth Newton-CG augmented Lagrangian method based on the software package SDPNAL+ [40]. Finally, numerical experiments on the QAPLIB [23] and ‘dre’ instances [18] demonstrate the proposed approach usually performs well.

Below are some common notations to be used in this paper. We use \mathcal{S}^q to denote the linear subspace of all $q \times q$ real symmetric matrices and use $\|\cdot\|$ to denote the Frobenius norm of matrices. Let $\mathcal{N}^q \subseteq \mathcal{S}^q$ be the subset of all $q \times q$ nonnegative symmetric matrices in \mathcal{S}^q . Denote $\mathcal{S}_+^q/\mathcal{S}_-^q$ ($\mathcal{S}_{++}^q/\mathcal{S}_{--}^q$) the positive/negative semidefinite (definite) matrix cone in \mathcal{S}^q . Moreover, let \mathcal{C}^q be the set of copositive matrices in \mathcal{S}^q and $(\mathcal{C}^q)^*$ be the dual cone of \mathcal{C}^q , i.e., the set of all completely positive matrices in \mathcal{S}^q . For a given matrix $Z \in \mathcal{S}^q$ with $q \geq 1$, we also use the following block notation for simplicity:

$$Z = \begin{bmatrix} Z^{11} & \dots & Z^{1q} \\ \vdots & \ddots & \vdots \\ Z^{q1} & \dots & Z^{qq} \end{bmatrix}$$

with $Z^{ij} \in \mathcal{R}^{q \times q}$ for each $i, j \in \{1, \dots, q\}$. Let e_i be the i -th standard unit vector. We denote the vector and square matrix of all ones by $\mathbf{1}_q$ and E_q respectively, and denote the identity matrix by I_q . We will omit the superscript q if the dimension is clear. For a given $Z \in \mathcal{S}^q$, we use $\lambda_1(Z) \geq \dots \geq \lambda_q(Z)$ to denote the eigenvalues of Z (all real and counting multiplicity) arranging in non-increasing order. We use “vec(\cdot)” to denote the vectorization of matrices and use “mat(\cdot)” to denote its inverse operator, i.e., the corresponding matrixization of vectors. If $z \in \mathcal{R}^q$, then $\text{Diag}(z)$ is a $q \times q$ diagonal matrix with z

on the main diagonal. Finally, we use “ \otimes ” to denote the Kronecker product between matrices.

2 The rank constrained DNN reformulation of the QAP

It is well-known that each permutation $\pi \in \mathcal{P}^n$ can be represented by a $n \times n$ permutation matrix X , i.e., a square binary matrix which has exactly one entry of 1 in each row and each column and zeros elsewhere. Therefore, the QAP (1) can be reformulate as the following trace form:

$$\min \{ \langle X, AXB + C \rangle \mid X \in \Pi^{n \times n} \}, \quad (2)$$

where $\langle \cdot, \cdot \rangle$ stands for the standard trace inner product of matrices, i.e., $\langle Y, Z \rangle = \text{tr}(X^T Y)$ for $X, Y \in \mathcal{R}^{m \times n}$, and $\Pi^{n \times n}$ is the set of all $n \times n$ permutation matrices. It is clear that $\Pi^{n \times n}$ can be characterized by the intersection of the set of orthogonal matrices and the set of nonnegative matrices, i.e.,

$$\Pi^{n \times n} = \{ X \in \mathcal{R}^{n \times n} \mid X^T X = I, X \geq 0 \}.$$

Without loss of generality, we may assume that the data matrices A, B, C in (1) are nonnegative, i.e., $A, B, C \in \mathcal{N}$. Inspired by [4], Povh and Rendl [35] suggested to consider the following convex completely positive conic relaxation of the QAP (2):

$$\begin{aligned} & \min \langle B \otimes A + \text{Diag}(c), Y \rangle \\ & \text{s.t. } \sum_{i=1}^n Y^{ii} = I, \quad \langle I, Y^{ij} \rangle = \delta_{ij}, \quad i, j \in \{1, \dots, n\}, \\ & \quad \langle E, Y \rangle = n^2, \quad Y \in (\mathcal{C}^{n^2})^*, \end{aligned} \quad (3)$$

where $c = \text{vec}(C)$ and $\delta_{ij} = 1$ if $i = j$ and $\delta_{ij} = 0$ otherwise for $i, j \in \{1, \dots, n\}$. It is clear that for any $n \times n$ permutation matrix $X \in \Pi^{n \times n}$,

$$Y = \text{vec}(X)\text{vec}(X)^T, \quad X \in \Pi^{n \times n}. \quad (4)$$

is a feasible solution of (3). Furthermore, Povh and Rendl [35] shown that the optimal value of (3) is actually equal the optimal value of QAP (2). Unfortunately, the completely positive cone constrain $Y \in (\mathcal{C}^{n^2})^*$ is computational intractable. A useful strategy to handle this is to approximate the cone $(\mathcal{C}^{n^2})^*$ from the outside, e.g., the cone of symmetric doubly nonnegative matrices $\mathcal{S}_+^{n^2} \cap \mathcal{N}^{n^2}$. Thus, we obtain the following relaxation of the QAP (2):

$$\begin{aligned} & \min \langle B \otimes A + \text{Diag}(c), Y \rangle \\ & \text{s.t. } \sum_{i=1}^n Y^{ii} = I, \quad \langle I, Y^{ij} \rangle = \delta_{ij}, \quad i, j \in \{1, \dots, n\}, \\ & \quad \langle E, Y \rangle = n^2, \quad Y \in \mathcal{S}_+^{n^2} \cap \mathcal{N}^{n^2}. \end{aligned} \quad (5)$$

Clearly, the optimal value of problem (5) only provides a lower bound of the QAP (2). In general, the relaxation (5) for the QAP is not tight.

On the other hand, from the equation (4), we may add the rank constraint $\text{rank}(Y) \leq 1$ to (5) and obtain the following rank constrained doubly nonnegative (DNN) problem:

$$\begin{aligned} & \min \langle B \otimes A + \text{Diag}(c), Y \rangle \\ & \text{s.t. } \sum_{i=1}^n Y^{ii} = I, \quad \langle I, Y^{ij} \rangle = \delta_{ij}, \quad i, j \in \{1, \dots, n\}, \\ & \quad \langle E, Y \rangle = n^2, \quad Y \in \mathcal{S}_+^{n^2} \cap \mathcal{N}^{n^2}, \quad \text{rank}(Y) \leq 1. \end{aligned} \quad (6)$$

The resulting problem (6) is non-convex. In fact, we shall show that (6) is an exact reformulation of the original QAP (2). To this end, we need the following simple observation on the rank one completely positive matrices.

Lemma 1 *Let $q \geq 1$ be a given positive integer. Suppose that $Y \in \mathcal{S}^q$ and $\text{rank}(Y) \leq 1$. Then, the following statements are equivalent:*

- (i) $Y \in (\mathcal{C}^q)^*$;
- (ii) $Y \in \mathcal{S}_+^q \cap \mathcal{N}^q$;
- (iii) there exists $x \in \mathcal{R}_+^q$ such that $Y = xx^T$.

Proof Since “(i) \implies (ii)” and “(iii) \implies (i)” are obvious, we only need to show “(ii) \implies (iii)”, i.e., if $Y \in \mathcal{S}_+^q \cap \mathcal{N}^q$, then there exists $x \in \mathcal{R}_+^q$ such that $Y = xx^T$. Without loss of generality, we may assume $\text{rank}(Y) = 1$, since otherwise the result holds trivially. It follows from $Y \in \mathcal{S}_+^q$ and $\text{rank}(Y) = 1$ that there exists $u \in \mathcal{R}^q$ such that $Y = \lambda uu^T$. Since $Y \geq 0$, we have $Y_{ij} = u_i u_j \geq 0$ for each $i, j \in \{1, \dots, q\}$. Thus, we can choose $x = \sqrt{\lambda}u \in \mathcal{R}_+^q$ such that $Y = xx^T$. \square

It is clear that the objective functions of (2) and (6) coincide. The equivalence between (2) and (6) then follows if we show the feasible sets of these two problems are the same. By employing the similar argument as that of [35, Theorem 3], we have the following result on the equivalence of the feasible sets of (6) and (2).

Proposition 1 *The matrix $Y \in \mathcal{S}_+^{n^2}$ is a feasible solution of (6) if and only if there exists a unique $X \in \Pi^{n \times n}$ such that $Y = \text{vec}(X)\text{vec}(X)^T$. Moreover, since $\|\text{vec}(X)\|^2$ is the only nonzero eigenvalue of Y , the vector $\text{vec}(X)/\|\text{vec}(X)\|$ is the unit nonnegative eigenvector of Y .*

Proof It is easy to see that if $X \in \Pi^{n \times n}$ then $Y = \text{vec}(X)\text{vec}(X)^T$ belongs the feasible set of (6). Thus, we only need to show the converse direction holds. Suppose that Y is a feasible set of (6). We know that $\text{rank}(Y) = 1$, since $Y \neq 0$. It then follows from Lemma 1 that there exists $y \in \mathcal{R}_+^{n^2}$ such that $Y = yy^T$ and $X = \text{mat}(y) \in \mathcal{R}^{n \times n}$. Then, by employing the similar argument as that of [35, Theorem 3], we are able to show that $X \in \Pi^{n \times n}$. Furthermore,

it is easy to verify that for any $X, X' \in \Pi^{n \times n}$, if $X \neq X'$, then $Y \neq Y'$ with $Y = \text{vec}(X)\text{vec}(X)^T$ and $Y' = \text{vec}(X')\text{vec}(X')^T$.

Let the nonzero unit vector $v \in \mathcal{R}^{n^2}$ with $v = \text{vec}(X)/\|\text{vec}(X)\|$, Obviously, $v \in \mathcal{R}_+^{n^2}$. From the definition of the characteristic polynomial for matrices, we know that

$$Yv = \text{vec}(X)\text{vec}(X)^T \cdot \frac{\text{vec}(X)}{\|\text{vec}(X)\|} = \|\text{vec}(X)\|\text{vec}(X) = \|\text{vec}(X)\|^2 v,$$

that is, $\|\text{vec}(X)\|^2$ and v is the eigenvalue and eigenvector of Y respectively. The proof is completed. \square

Remark 1 It follows from Proposition 1 that if $Y \in \mathcal{S}^{n^2}$ is a feasible solution of (6), then we can find the permutation matrix $X \in \Pi^{n \times n}$ by setting $X = \text{mat}(x)$ with $x = v \cdot \|\text{vec}(X)\|$ easily, where v is the unit corresponding eigenvector of Y with respect to n .

The following result on the equivalence between the rank constrained DNN problem (6) and the QAP (2) follows from Proposition 1 immediately.

Theorem 1 *The rank constrained DNN problem (6) is equivalent to the QAP (2).*

Clearly, the nonconvex rank constrained DNN representation (6) is at least as hard as the original QAP, which means that finding a global solution of (6) is computationally intractable. However, it is still possible to design some efficient algorithms, e.g., the DCA (see Section 4), to find a good feasible point of (6) and obtain a good feasible solution of the original QAP.

3 Extensions

In this section, we shall demonstrate that the results obtained in Section 2 can be applied to other important nonconvex problems, which have the similar rank constrained DNN representations.

Standard quadratic programming. The standard quadratic problem (StQP) consists of finding an optimal of a quadratic form over the standard simplex, i.e.,

$$\min \left\{ \langle x, Qx \rangle \mid \sum_{i=1}^n x_i = 1, x \geq 0 \right\}, \quad (7)$$

where Q is an arbitrary $n \times n$ symmetric matrix. The StQP (7) includes many important combinatorial optimization problems as special cases, e.g., the maximum clique problem [32]. It is clear that the StQP (7) can be rewritten as the following matrix form:

$$\begin{aligned} & \min \langle Q, Y \rangle \\ & \text{s.t. } \langle E, Y \rangle = 1, \quad Y = xx^T, \quad x \geq 0. \end{aligned}$$

Thus, by employing Lemma 1, we obtain the following result on the rank constrained DNN representation of the StQP (7), immediately.

Theorem 2 *The standard quadratic problem (7) is equivalent to the following rank constrained DNN problem:*

$$\begin{aligned} \min \langle Q, Y \rangle \\ \text{s.t. } \langle E, Y \rangle = 1, \quad Y \in \mathcal{S}_+^n \cap \mathcal{N}^n, \quad \text{rank}(Y) \leq 1. \end{aligned} \quad (8)$$

The minimum-cut graph tri-partitioning problem. The minimum-cut graph tri-partitioning problem [34] is to find a tri-partitioning the vertices of a graph into sets S_1 , S_2 and S_3 of specified cardinalities, such that the total weight of edges between S_1 and S_2 is minimal.

Let $G = (V, E)$ be an undirected graph on n vertices, given by its (weighted) symmetric nonnegative adjacency matrix $A \in \mathcal{N}^n$, the minimum-cut graph tri-partitioning problem [34] can be described as: for given integers m_1 , m_2 and m_3 summing to n , find subsets S_1 , S_2 and S_3 of $V(G)$ with cardinalities m_1 , m_2 and m_3 , respectively, such that the total weight of edges between S_1 and S_2 is minimal. By presenting partitions S_1 , S_2 and S_3 by $n \times 3$ matrices X , the minimum-cut graph tri-partitioning problem can be written as follows

$$\begin{aligned} \min \frac{1}{2} \langle X, AXB \rangle \\ \text{s.t. } X^T X = M, \quad X \mathbf{1}_3 = \mathbf{1}_n, \\ X \geq 0, \end{aligned} \quad (9)$$

where $M := \text{Diag}(m_1, m_2, m_3)$ and $B = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, the vector of all ones is

$\mathbf{1}_k \in \mathcal{R}^k$. By introducing $Y = xx^T$ with $x = \text{vec}(X)$, Povh and Rendl [34] reformulate the minimum-cut graph tri-partitioning problem (9) as follows:

$$\begin{aligned} \min \frac{1}{2} \langle B \otimes A, Y \rangle \\ \text{s.t. } \langle L^{ij} \otimes I, Y \rangle = m_i \delta_{ij}, \quad 1 \leq i \leq j \leq 3, \\ \langle E_3 \otimes J^{ii}, Y \rangle = 1, \quad 1 \leq i \leq n, \\ \langle V_i \otimes W_j^T, Y \rangle = m_i, \quad 1 \leq i \leq 3, 1 \leq j \leq n \\ \langle L^{ij} \otimes E_n, Y \rangle = m_i m_j, \quad 1 \leq i \leq j \leq 3, \\ Y = xx^T, x \in \mathcal{R}_+^{3n}, \end{aligned} \quad (10)$$

where $V_i = e_i \mathbf{1}_3^T \in \mathcal{R}^{3 \times 3}$ for $i = 1, 2, 3$, $W_j = e_j \mathbf{1}_n^T \in \mathcal{R}^{n \times n}$ for $j = 1, \dots, n$, $J^{ij} = e_i e_j^T \in \mathcal{R}^{n \times n}$ and $L^{ij} = \frac{1}{2}(e_i e_j^T + e_j e_i^T) \in \mathcal{R}^{3 \times 3}$ for $i, j = 1, 2, 3$. Again, similar with Section 2, by employing Lemma 1, we are able to obtain the following rank constrained DNN representation of the minimum-cut graph tri-partitioning problem (9).

Theorem 3 *The minimum-cut graph tri-partitioning problem (9) is equivalent to the following rank constrained DNN problem:*

$$\begin{aligned}
& \min \frac{1}{2} \langle B \otimes A, Y \rangle \\
& \text{s.t. } \langle L^{ij} \otimes I, Y \rangle = m_i \delta_{ij}, \quad 1 \leq i \leq j \leq 3, \\
& \quad \langle E_3 \otimes J^{ii}, Y \rangle = 1, \quad 1 \leq i \leq n, \\
& \quad \langle V_i \otimes W_j^T, Y \rangle = m_i, \quad 1 \leq i \leq 3, 1 \leq j \leq n \\
& \quad \langle L^{ij} \otimes E_n, Y \rangle = m_i m_j, \quad 1 \leq i \leq j \leq 3, \\
& \quad Y \in \mathcal{S}_+^{3n} \cap \mathcal{N}^{3n}, \quad \text{rank}(Y) \leq 1.
\end{aligned} \tag{11}$$

4 The DCA for the rank constrained DNN problem

In this section, we shall propose a DCA based algorithm for the rank constrained DNN relaxations established in the previous section. For simplicity in notation, all proposed rank constrained DNN representations (6), (8) and (11) can be cast in the following abstract form:

$$\begin{aligned}
& \min f(Y) := \langle \bar{C}, Y \rangle \\
& \text{s.t. } Y \in \Omega \cap \mathcal{R},
\end{aligned} \tag{12}$$

where the subsets $\Omega, \mathcal{R} \subseteq \mathcal{S}^q$ are defined by

$$\Omega := \left\{ Y \in \mathcal{S}_+^q \cap \mathcal{N}^q \mid \mathcal{A}(Y) = b \right\} \tag{13}$$

and

$$\mathcal{R} := \{ Y \in \mathcal{S}^q \mid \text{rank}(Y) \leq 1 \}, \tag{14}$$

$\bar{C} \in \mathcal{S}^q$, $\mathcal{A} : \mathcal{S}^q \rightarrow \mathcal{R}^m$ is a given linear operator, and $b \in \mathcal{R}^m$ is a given data.

It is worth to note that for the rank constrained DNN relaxations proposed in Section 2, the subsets Ω with respect to (6), (8) and (11) [satisfy the](#) following assumption.

Assumption 1 *The subset $\Omega \subseteq \mathcal{S}^q$ defined by (13) is nonempty and bounded.*

It is well-known that for any $Y \in \mathcal{S}_+^q$, $\text{rank}(Y) \leq 1$ if and only if $0 = \lambda_2(Y) + \dots + \lambda_n(Y) = \sum_{i=1}^n \lambda_i(Y) - \lambda_1(Y) = \|Y\|_* - \|Y\|_2$, where $\|\cdot\|_*$ and $\|\cdot\|_2$ denote the nuclear norm (i.e., the sum of all singular values) and spectral norm (i.e., the largest singular value) of matrices, respectively. Therefore, the rank constraint DNN problem (12) can be rewritten as follows:

$$\begin{aligned}
& \min f(Y) \\
& \text{s.t. } \|Y\|_* - \|Y\|_2 = 0 \\
& \quad Y \in \Omega.
\end{aligned} \tag{15}$$

Furthermore, by penalizing the second constraint in (15) into the objective function, we obtain the following penalized problem

$$\begin{aligned} \min f_\rho(Y) &:= \langle \bar{C}, Y \rangle + \rho(\|Y\|_* - \|Y\|_2) \\ \text{s.t. } Y &\in \Omega, \end{aligned} \quad (16)$$

where $\rho > 0$ is a given penalty parameter.

In order to establish the exact penalty property of (16) with respect to (15), we need further introduce the following concept on the calmness [16] of set-valued mappings. Let \mathcal{X} and \mathcal{Z} be two finite dimensional Euclidean space. Recall a set-valued mapping $\Psi : \mathcal{X} \rightrightarrows \mathcal{Z}$ is called calm at \bar{x} for $\bar{z} \in \Psi(\bar{x})$ if there exist a constant $\alpha > 0$ and neighborhood $\mathcal{U} \subseteq \mathcal{X}$ of \bar{x} and neighborhood $\mathcal{V} \subseteq \mathcal{Z}$ of \bar{z} such that

$$\Psi(x) \cap \mathcal{V} \subseteq \Psi(\bar{x}) + \alpha\|x - \bar{x}\|\mathbb{B}_{\mathcal{X}} \quad \forall x \in \mathcal{U},$$

where $\mathbb{B}_{\mathcal{X}}$ is the unit ball in \mathcal{X} .

Proposition 2 *Suppose that the set-valued mapping $\Gamma : \Re \rightrightarrows \mathcal{S}^q$ defined by*

$$\Gamma(w) := \{Y \in \mathcal{S}^q \mid Y \in \Omega, \|Y\|_* - \|Y\|_2 = w\}, \quad w \in \Re, \quad (17)$$

is calm at 0 for each $Y \in \Gamma(0)$. Then, there exists a constant $\bar{\rho} > 0$ such that for any $\rho > \bar{\rho}$, Y^ is an optimal of (12) if and only if Y^* is an optimal of the penalized problem (16).*

Proof First, we shall show that there exists $\bar{\rho} > 0$ if Y^* is an optimal of (12), then it is also an optimal solution of the penalized problem (16) for $\rho > \bar{\rho}$. By [8, Theorem 2.1], we know from the calmness of Γ that there exists $\tau > 0$ such that $\text{dist}(Y, \Omega \cap \mathcal{R}) \leq \tau \text{dist}(Y, \mathcal{R}) = \tau(\|Y\|_* - \|Y\|_2)$ for any $Y \in \Omega$. Let $L := \|\bar{C}\| > 0$. Suppose that $\rho > \bar{\rho} := \max\{L\tau, L\}$ be arbitrarily given. Suppose there exists $\bar{Y} \in \Omega$ and $\varepsilon > 0$ such that

$$\langle \bar{C}, \bar{Y} \rangle + \rho(\|\bar{Y}\|_* - \|\bar{Y}\|_2) < \langle \bar{C}, Y^* \rangle - \rho\varepsilon.$$

Let $\hat{Z} \in \Omega \cap \mathcal{R}$ be such that

$$\|\hat{Z} - \bar{Y}\| \leq \text{dist}(\bar{Y}, \Omega \cap \mathcal{R}) + \varepsilon.$$

Since $\text{dist}(\bar{Y}, \Omega \cap \mathcal{R}) \leq \tau \text{dist}(\bar{Y}, \mathcal{R}) = \tau(\|\bar{Y}\|_* - \|\bar{Y}\|_2)$, we have

$$\|\hat{Z} - \bar{Y}\| \leq \tau(\|\bar{Y}\|_* - \|\bar{Y}\|_2) + \varepsilon.$$

Then,

$$\begin{aligned} \langle \bar{C}, \hat{Z} \rangle &\leq \langle \bar{C}, \bar{Y} \rangle + L\|\hat{Z} - \bar{Y}\| \leq \langle \bar{C}, \bar{Y} \rangle + L(\tau(\|\bar{Y}\|_* - \|\bar{Y}\|_2) + \varepsilon) \\ &\leq \langle \bar{C}, \bar{Y} \rangle + \rho(\|\bar{Y}\|_* - \|\bar{Y}\|_2 + \varepsilon) < \langle \bar{C}, Y^* \rangle. \end{aligned}$$

This contradicts with the fact that Y^* is an optimal of (12).

For the converse direction, it is sufficient to show that if Y^* is an optimal of the penalized problem (16), then $Y^* \in \Omega \cap \mathcal{R}$, i.e., Y^* is a feasible solution of (12). In fact, if $\tilde{Y} \in \Omega \cap \mathcal{R}$ is an optimal of (12), then since Y^* is an optimal of the problem (16), we know from the first part that

$$\langle \bar{C}, Y^* \rangle + \rho(\|Y^*\|_* - \|Y^*\|_2) = \langle \bar{C}, \tilde{Y} \rangle$$

and

$$\langle \bar{C}, Y^* \rangle + \frac{1}{2}(\rho + \bar{\rho})(\|Y^*\|_* - \|Y^*\|_2) \geq \langle \bar{C}, \tilde{Y} \rangle,$$

which implies that

$$\frac{1}{2}(\bar{\rho} - \rho)(\|Y^*\|_* - \|Y^*\|_2) \geq 0.$$

Since $\rho > \bar{\rho}$ and $\|Y^*\|_* - \|Y^*\|_2 \geq 0$, we know that $\|Y^*\|_* - \|Y^*\|_2 = 0$, i.e., $\text{rank}(Y^*) \leq 1$. Thus, we have $Y^* \in \Omega \cap \mathcal{R}$. This completes the proof. \square

The calmness assumption on the set-valued mapping Γ defined by (17) may be difficult to verified for the general rank constrained DNN problem (12) including the models (6), (8) and (11) introduced in this paper. However, the calmness assumption is indeed satisfied for some classes of rank constraint SDP problems. See [8] for more details.

It is clear that the non-convex objective function in (16) is a DC (difference of convex) function. Therefore, the DC based method [41, 43] can be employed for solving (16). In particular, by employing the fact that $\|Y\|_* = \langle I, Y \rangle$ for any $Y \in \mathcal{S}_+^q$, we rewrite the objective function in (16) as the following special DC form:

$$f_\rho(Y) = \underbrace{\left(\langle \bar{C}, Y \rangle + \rho \langle I, Y \rangle + \frac{1}{2\sigma} \|Y\|^2 \right)}_{\text{denoted as } g(Y)} - \underbrace{\left(\rho \|Y\|_2 + \frac{1}{2\sigma} \|Y\|^2 \right)}_{\text{denoted as } h(Y)}, \quad (18)$$

where $\sigma > 0$ is a given parameter. The regularization term $\frac{1}{2\sigma} \|\cdot\|^2$ is widely used in DCA (see e.g., [42, (16)] for the application in solving trust region subproblems) and well-studied in literature (cf.e.g., [41, Section 5] for details).

Next, we shall employ the simplified DCA introduced by [41] to solve (16) based on the DC regularized decomposition (18). Let $Y^0 \in \Omega$ be given. For $k \geq 0$, at k -th iteration, the simplified DCA then takes the following form: let $Z^k := \rho W^k + \frac{1}{\sigma} Y^k$ with $W^k \in \partial \|Y^k\|_2$ and Y^{k+1} be the optimal solution of the following strongly convex problem:

$$\begin{aligned} & \min g(Y) - \langle Y, Z^k \rangle \\ & \text{s.t. } Y \in \Omega. \end{aligned} \quad (19)$$

It is clear that (19) is equivalent to the following problem:

$$\begin{aligned} & \min \langle \bar{C}, Y \rangle + \rho \langle I - W^k, Y \rangle + \frac{1}{2\sigma} \|Y - Y^k\|^2 \\ & \text{s.t. } Y \in \Omega. \end{aligned} \quad (20)$$

Thus, the simplified DCA for solving (16) based on the decomposition (18) takes the following form.

Algorithm 1 [Proximal DC Algorithm (ProxDCA)]

- 1: Let $Y^0 \in \Omega$ be an initial point and $\sigma > 0$. Set $k = 0$.
 2: Compute

$$Y^{k+1} = \operatorname{argmin} \left\{ \hat{f}_\rho^k(Y) \mid Y \in \Omega \right\}, \quad (21)$$

where $\hat{f}_\rho^k(Y) := \langle \bar{C}, Y \rangle + \rho \langle I - W^k, Y \rangle + \frac{1}{2\sigma} \|Y - Y^k\|^2$ and $\Omega \subseteq \mathcal{S}^q$ is defined by (13).

- 3: If $Y^{k+1} = Y^k$ stop; otherwise set $k = k + 1$, go to **Step.2**.
-

Under Assumption 1, the strongly convex problem (21) has a unique solution and can be solved efficiently by considering its dual problem, i.e.,

$$\begin{aligned} \max \quad & -\langle b, y \rangle - \frac{\sigma}{2} \left\| \bar{C} + \rho(I - W^k) + \mathcal{A}^*y + S + Z - \frac{1}{\sigma}Y^k \right\|^2 \\ \text{s.t.} \quad & S \in \mathcal{S}_-^q, \quad Z \in -\mathcal{N}^q, \end{aligned} \quad (22)$$

If $(y^{k+1}, S^{k+1}, Z^{k+1}) \in \mathcal{R}^m \times \mathcal{S}^q \times \mathcal{N}^q$ is an optimal solution of the above dual problem (22), then Y^{k+1} can be found as follows

$$Y^{k+1} = Y^k - \sigma (\mathcal{A}^*y^{k+1} + S^{k+1} + Z^{k+1} + \bar{C} + \rho(I - W^k)).$$

Moreover, it is clear that the dual problem (22) is equivalent to the augmented Lagrangian sub-problem [48, (7a)], which can be solved efficiently by employing the majorized semismooth Newton-CG algorithm **Algorithm MSNCG** introduced by [48]. Furthermore, under the following general Slater condition, the convergence of **Algorithm MSNCG** is established in [48, Theorem 2.5].

Assumption 2 *There exists $\tilde{Y} \in \mathcal{S}^q$ such that*

$$\mathcal{A}(\mathcal{T}_{\mathcal{N}^q}(\tilde{Y})) = \mathcal{R}^m \quad \text{and} \quad \tilde{Y} \in \mathcal{S}_{++}^q \cap \operatorname{int}(\mathcal{N}^q),$$

where $\operatorname{int}(\mathcal{N}^q)$ and $\mathcal{T}_{\mathcal{N}^q}(\tilde{Y})$ denote the interior of \mathcal{N}^q and the tangent cone of \mathcal{N}^q at \tilde{Y} , respectively.

Note that for the constraint set Ω defined by (13), $\mathcal{T}_{\mathcal{N}^q}(\tilde{Y})$ is actually the space \mathcal{S}^q as \tilde{Y} is assumed to be in the interior of \mathcal{N}^q . For the proposed rank constrained DNN models e.g., (6), (8) and (11), Assumption 2 will be satisfied since the linear mappings \mathcal{A} in these models are usually full row rank.

Next, we shall study the convergence of the proposed DCA based algorithm for the rank constrained DNN problem (12). A feasible point $Y \in \Omega$ is said to be a stationary point of the penalized problem (16) if

$$(\bar{C} + \rho I + \mathcal{N}_\Omega(Y)) \cap (\rho \partial \|Y\|_2) \neq \emptyset,$$

where $\mathcal{N}_\Omega(Y)$ is the normal cone of the convex set Ω at Y in the sense of convex analysis (cf. e.g., [38]). The following results on the convergence of the proposed DC based algorithm (Algorithm 1) for the rank constrained DNN problem (12) follows from the basic convergence theorem of DCA [41, Theorem 3] immediately. For simplicity, we omit the detail proof here.

Proposition 3 *Suppose that Assumption 1 holds. Let $\rho > 0$ be given. Let $\{Y^k\}$ be the sequence generated by Algorithm 1. Then $\{f_\rho(Y^k)\}$ is a monotonically decreasing sequence. If $Y^{k+1} = Y^k$ for some integer $k \geq 0$, then Y^{k+1} is a stationary point of the penalized problem (16). Otherwise, the infinite sequence $\{f_\rho(Y^k)\}$ satisfies*

$$\frac{1}{\sigma} \|Y^{k+1} - Y^k\|^2 \leq f_\rho(Y^k) - f_\rho(Y^{k+1}), \quad k = 0, 1, \dots \quad (23)$$

Moreover, any accumulation point of the bounded sequence $\{Y^k\}$ is a stationary point of problem (16).

In order to show the infinity sequence $\{Y^k\}$ generated by the proposed Algorithm 1 actually converge, we recall the following definition of the Kurdyka-Lojziewicz (KL) property of the lower semi-continuous function (see [5, 11, 12] for more details). Let $\iota > 0$ and Ψ_ι be the class of functions $\psi : [0, \iota) \rightarrow \mathcal{R}_+$ that satisfy the following conditions:

- (a) $\psi(0) = 0$;
- (b) ψ is positive, concave and continuous;
- (c) ψ is continuously differentiable on $(0, \iota)$ with $\psi'(x) > 0$ for any $x \in (0, \iota)$.

Let $g : \mathcal{R}^n \rightarrow (-\infty, \infty]$ be a given proper lower semicontinuous function. Suppose that $x \in \text{dom } g := \{x \in \mathcal{R}^n \mid g(x) < \infty\}$. The Fréchet subdifferential of g at x is defined as

$$\widehat{\partial} g(x) := \left\{ h \in \mathcal{R}^n \mid \limsup_{x \neq y \rightarrow x} \frac{g(y) - g(x) - h^T(y - x)}{\|y - x\|} \geq 0 \right\}$$

and the limiting subdifferential, or simply the subdifferential of g at x , is defined by

$$\partial g(x) := \left\{ h \in \mathcal{R}^n \mid \exists \{x^k\} \rightarrow x \text{ and } \{h^k\} \rightarrow h \text{ satisfying } h^k \in \widehat{\partial} g(x^k) \forall k \right\}.$$

Definition 1 (KL property) The given proper lower semicontinuous function $g : \mathcal{R}^n \rightarrow (-\infty, \infty]$ is said to have the KL property at $\bar{x} \in \text{dom } g$ if there exist $\iota > 0$, a neighborhood \mathcal{U} of \bar{x} and a concave function $\psi \in \Psi_\iota$ such that

$$\psi'(g(x) - g(\bar{x})) \text{dist}(0, \partial g(x)) \geq 1 \quad \forall x \in \mathcal{U} \text{ and } g(\bar{x}) < g(x) < g(\bar{x}) + \iota,$$

where $\text{dist}(x, Z) = \min_{z \in Z} \|y - x\|$ is the distance from a point x to a nonempty closed set Z . The function g is said to be a KL function if it has the KL property at each point of $\text{dom } g$.

One most frequently used functions which has the KL property are the semialgebraic functions.

Definition 2 (Semialgebraic sets and functions) A set in \mathcal{R}^n is *semialgebraic* if it is a finite union of sets of the form:

$$\{x \in \mathcal{R}^n \mid p_i(x) > 0, q_j(x) = 0, \quad i = 1, \dots, a, j = 1, \dots, b\},$$

where $p_i : \mathcal{R}^n \rightarrow \mathcal{R}$, $i = 1, \dots, a$ and $q_j : \mathcal{R}^n \rightarrow \mathcal{R}$, $j = 1, \dots, b$ are polynomials. A mapping is semialgebraic if its graph is semialgebraic.

For this class of function, we have the following useful result (cf. [9, 10]).

Proposition 4 *Suppose a proper lower semicontinuous function $g : \mathcal{R}^n \rightarrow (-\infty, \infty]$ is semialgebraic, then g is a KL function.*

It is well-known that a symmetric matrix is positive semidefinite if and only if all its principal minors are non-negative. Thus, the positive semidefinite matrix cone can be written as the intersection of finite many polynomial inequalities, which is indeed a semialgebraic set. Moreover, the semialgebraic property is stable under the boolean operations, i.e., finite unions and intersections and taking complement (cf. e.g., [19]). We know that the set Ω is semialgebraic. Moreover, it is known from the Tarski-Seidenberg theorem (see e.g., [31, Theorem 8.6.6]), the semidefinite programming representable sets [7, Section 4.2] are all semialgebraic (cf. e.g., [25] for more details).

Now, we are ready to establish the global convergence of Algorithm 1 by employing a refined global convergence result for the proximal DCA solving the DC programming with the nonsmooth DC function, which is recently developed by Liu et al. [30].

Theorem 4 *Suppose that Assumption 1 holds. Let $\rho > 0$ be given and $0 < \sigma \leq 1/\|\bar{C} + \rho I\|$. Suppose that $\{Y^k\}$ is the infinite sequence generated by Algorithm 1. Then $\{Y^k\}$ converges to a stationary point of problem (16).*

Proof As mentioned before, the set $\Omega \subseteq \mathcal{S}^q$ defined in (13) is semialgebraic. On the other hand, the conjugate function $p^*(Y) := \sup_{Z \in \mathcal{S}^q} \{\langle Y, Z \rangle - \|Z\|_2\}$ coincides with the indicator function of the unit ball of the nuclear norm $\|\cdot\|_*$, i.e., $\{Y \in \mathcal{S}^n \mid \|Y\|_* \leq 1\}$ (cf. [38, Theorems 13.5 & 13.2]). It is well-known [7, Example 18(c) & 19] that the unit ball of the nuclear norm is semidefinite programming representable. Thus, we know that the conjugate function $p^*(Y)$ is semialgebraic. Moreover, by the fact that finite sums of semialgebraic functions are semialgebraic (see, e.g., [6, Section 4.3]), we know that given $\sigma > 0$, the corresponding auxiliary major function $E(Y, Z, W) := \langle \bar{C}, Y \rangle + \rho \langle I, Y \rangle + \delta_\Omega(Y) - \langle Y, Z \rangle + p^*(Z) + \frac{1}{2\sigma} \|Y - W\|^2$, $Y, Z, W \in \mathcal{S}^n$ defined in [30, (7)] is semialgebraic. It then follows from Proposition 4 that the function E is a KL function. Furthermore, it can be seen from [30, Remark 3.1] that Algorithm 1 is a special variant of the proximal difference-of-convex algorithm with extrapolation (pDCA_e) proposed by Liu et al. [30] (with $\beta_k \equiv 0$ in [30, (5)]). Thus, the desired result follows from [30, Theorem 3.1] directly. \square

Finally, we will show that if the parameter $\rho > 0$ is large enough, then the sequence $\{Y^k\}$ obtained by Algorithm 1 will satisfy the rank constraint of (12) when k sufficiently large.

Proposition 5 *Suppose that Assumptions 1 and 2 hold. For each k , choose $W^k = U_1^k (U_1^k)^T \in \partial p(Y^k)$, where $U_1^k \in \mathcal{R}^q$ is the orthonormal eigenvector with respect to the largest eigenvalue $\lambda_1(Y^k)$ of Y^k . Let $\{Y^k\}$ be the sequence generated by Algorithm 1 and $\{y^k, S^k, Z^k\}$ be the corresponding dual solution sequence of (22). If there exists a finite constant η such that for k sufficiently large,*

$$\lambda_1(-\bar{C} - \mathcal{A}^* y^{k+1} - Z^{k+1}) \leq \eta, \quad (24)$$

then there exists $\hat{\rho} > 0$ such that for any $\rho > \hat{\rho}$ and each k sufficiently large,

$$\text{rank}(Y^{k+1}) \leq 1,$$

which implies that Y^{k+1} is a feasible solution of (12).

Proof For each k , since problem (21) is convex, we know that under Assumption 1 and 2, $Y^{k+1} \in \mathcal{S}^q$ and $(y^{k+1}, S^{k+1}, Z^{k+1}) \in \mathcal{R}^m \times \mathcal{S}^q \times \mathcal{S}^q$ satisfy the following KKT system:

$$\begin{cases} \bar{C} + \rho(I - W^k) + \mathcal{A}^* y^{k+1} + S^{k+1} + Z^{k+1} + \frac{1}{\sigma}(Y^{k+1} - Y^k) = 0, \\ \mathcal{A}(Y^{k+1}) = b, \\ \mathcal{S}_+^q \ni Y^{k+1} \perp S^{k+1} \in \mathcal{S}_-^q, \quad \mathcal{N}^q \ni Y^{k+1} \perp Z^{k+1} \in -\mathcal{N}^q. \end{cases} \quad (25)$$

By the first equation of (25), we know that for each k ,

$$S^{k+1} = -\rho(I - W^k) + M^{k+1},$$

where $M^{k+1} := -\bar{C} - \mathcal{A}^* y^{k+1} - Z^{k+1} - \frac{1}{\sigma}(Y^{k+1} - Y^k)$. By Weyl's eigenvalue inequality (see [47] or [24, Theorem 4.3.7]), we have for each k ,

$$\lambda_2(S^{k+1}) \leq \lambda_2(-\rho(I - W^k)) + \lambda_1(M^{k+1}) = -\rho + \lambda_1(M^{k+1}), \quad (26)$$

where the equality holds due to the fact that the eigenvalues $\lambda(-\rho(I - W^k)) = (0, -\rho, \dots, -\rho)^T \in \mathcal{R}^q$. Moreover, since for each k , $Y^k \in \Omega$ is bounded, we know that there exists a constant $\zeta > 0$ such that for each k , $\|Y^k\|_2 \leq \zeta$. Since there exists a finite constant η such that for k sufficiently large, $\lambda_1(-\bar{C} - \mathcal{A}^* y^{k+1} - Z^{k+1}) \leq \eta$, we know that there exists a constant $\zeta > 0$ such that for k sufficiently large,

$$\begin{aligned} \lambda_1(M^{k+1}) &\leq \lambda_1(-\bar{C} - \mathcal{A}^* y^{k+1} - Z^{k+1}) + \frac{1}{\sigma} \lambda_1(Y^{k+1} - Y^k) \\ &\leq \lambda_1(-\bar{C} - \mathcal{A}^* y^{k+1} - Z^{k+1}) + \frac{1}{\sigma} \|Y^{k+1} - Y^k\|_2 \leq \eta + \frac{\zeta}{\sigma}. \end{aligned}$$

Therefore, we know from (26) that if $\rho > \hat{\rho} := \max\{\eta, 0\} + \frac{\zeta}{\sigma} > 0$, then for k sufficiently large,

$$\lambda_2(S^{k+1}) \leq -\rho + \lambda_1(M^{k+1}) \leq -\rho + \eta + \frac{\zeta}{\sigma} < 0. \quad (27)$$

Finally, since $\mathcal{S}_+^q \ni Y^{k+1} \perp S^{k+1} \in \mathcal{S}_-^q$, by (27), we obtain that for k sufficiently large,

$$\text{rank}(S^{k+1}) \geq q - 1 \quad \text{and} \quad \text{rank}(Y^{k+1}) + \text{rank}(S^{k+1}) \leq q,$$

which implies that $\text{rank}(Y^{k+1}) \leq q - \text{rank}(S^{k+1}) \leq 1$. \square

It seems difficult to show that for the rank constrained DNN problem (12), the dual solution sequence $\{y^k, S^k, Z^k\}$ of (22) satisfies (24) for some constant η . However, we will demonstrate few numerical examples that (24) is indeed satisfied in some cases.

5 Numerical results

In this section, we present numerical results for the relaxation problem (6) solving by Algorithm 1. All the data from QAPLIB [23] and ‘dre’ instances [18] are tested on a Window 10 workstation (6 core, Intel Xeon E5-2650 v3 @ 2.30 GHz, 128 GB RAM). The size of most QAPs ranges from 12 to 60. During our experiments, SDPNAL+ version 1.0 [40] is used as doubly non-negative solver for solving the subproblems (21). Algorithm 1 is implemented in the MATLAB 2015a platform. In numerical experiments, we choose the regularization parameter of the subproblem (21) as $\sigma = 0.9/\|\bar{C} + \rho I\|$. We measure the performance of Algorithm 1 by

$$\text{gap} := \frac{\text{PDCA} - \text{opt}}{\text{opt}} \times 100\%,$$

where ‘opt’ denotes the optimal value (or best-known feasible solution) of each instance from QAPLIB, ‘PDCA’ denotes the object function value of the problem (12) obtained by Algorithm 1.

5.1 Penalty parameter

The penalty parameter ρ is an important factor for the whole procedure of Algorithm 1. It is worth to note that for sufficiently large ρ , Algorithm 1 always produces a rank-one solution, i.e., a feasible solution of the rank constraint DNN problem (12). In fact, Figure 1 shows that the largest eigenvalue assumption in Proposition 5 for the dual solution sequence for different ρ is usually satisfied for the QAP problems: chr18a, els19, had20 and lipa30a. In fact, it can be seen from Figure 1 that for large k , the largest eigenvalue of matrix $-\bar{C} - \mathcal{A}^*y^{k+1} - Z^{k+1}$ usually closes to a constant or decreasing with

respect to the penalty parameter ρ . Figure 2 shows the effect of the parameter ρ on the gaps and the ranks of the sequences generated by Algorithm 1 for chr18a, els19, had20 and lipa30a. In each subfigure, the x-axis is the range of the parameter ρ , the left, and right y-axis denote the ranks of the generated solutions (the blue lines) and the gaps of the optimality (the red lines) for the different ρ respectively. As shown in Fig. 2 (a) and (b), if ρ increases from 0, chr18 and els19 problems can obtain the optimal solutions of the problem (12) since the gaps are zeroes.




Fig. 1: Effects of parameters ρ on ranks and the largest eigenvalues of $-\bar{C} - \mathcal{A}^*y^{k+1} - Z^{k+1}$ for k sufficiently large.

Although larger ρ can help the solutions satisfying the rank-one constraint in the problem (12) (Proposition 5), the parameter ρ should not be too large. It is worth to know that the penalty problem (16) is close to the target problem (12) in the sense of the globe optimality for large ρ . However, the sequence $\{Y^k\}$ generated by Algorithm 1 only converges to a stationary point of (16) for a fixed ρ . When ρ increases, in the current stage, we still do not know whether the resulting stationary point is close to a global solution of (16) or not. As demonstrated by (c) had20 and (d) lipa30a in Fig. 2, the gaps between the objective function values obtained by Algorithm 1 and the known




Fig. 2: Effects of parameters ρ on gaps and ranks of solutions

global optimal values oscillate up and down when ρ increases. Therefore, in our implementation, a bisection strategy in $[0.001, 0.02]$ is used for finding a suitable parameter ρ for Algorithm 1.

5.2 Numerical performance

Table 1 summarizes the quality of the solutions obtained by our proposed ProxDCA approach for solving the problems from QAPLIB [23] and ‘dre’ instances [18] (107 instances). It can be seen from Table 1 that for 69 instances we are able to solve the problems exactly; for 32 instances we can obtain a feasible solution whose gap is less than or equal 4%; for 6 instances we obtain a feasible solution whose gap is larger than 4%.

The detail numerical results of Algorithm 1 for solving the ‘dre’ instances from [18] and QAPLIB [23] are reported in Tables 2 and 3. In these tables, ‘time’ column (in hours:minutes:seconds) reports the CPU time of Algorithm 1 and ‘permutation/bound’ column reports the feasible solution generated by

Table 1: Summary of numerical performance of Algorithm 1

Problem set (No.)	gap			Problem
	0	$\leq 4\%$	$> 4\%$	
drexxx(6)	6	0	0	dre15, dre18, dre21, dre24, dre30, dre42
bur26x(8)	0	8	0	bur26a-h
chrxxx(14)	14	0	0	chr12x, chr15x, chr18x, chr20x, chr22x, chr25a
els19(1)	1	0	0	els19
escxxx(14)	11	1	2	esc16a-j, esc32a-g
hadxx(5)	5	0	0	had12, had14-had20
kra32x(3)	1	2	0	kra30a-b, kra32
lipaxxx(10)	10	0	0	lipa20x, lipa30x, lipa40x, lipa50x, lipa60x
nugxx(13)	8	5	0	nug12, nug14-nug22, nug25, nug27, nug28
rouxx(3)	3	0	0	rou12, rou15, rou20
scrxx(3)	3	0	0	scr12, scr15, scr20
skoxx(5)	0	2	3	sko42, sko56, sko64, sko72, sko81
ste36x(3)	0	3	0	ste36a-c
taixxx(17)	7	9	1	tai12x, tai15x, tai17x, tai20x, tai25x, tai30x, tai35x, tai40x, tai50a, tai60b
thoxx(2)	0	2	0	tho30, tho40
Total(107)	69	32	6	

solving the relaxation problem (21) of the rank-1 constrained DNN problem (16).

The ‘dre’ problem instances [18] are based on a rectangular grid where all nonadjacent nodes have zero weight, making the value of the objective function increase steeply with just a slight change from the optimal permutation. The ‘dre’ instances are difficult to solve, especially for many metaheuristic-based methods, since they are ill-conditioned and hard to break out the ‘basin’ of the local minimal. The best-known solutions for the ‘dre’ problems have been found by branch and bound in [18]. Notably, by employing our proposed DCA based approach Algorithm 1, we are able to obtain the global optimal solutions of the ‘dre’ problems quite efficiently. For instance, we are able to solve the instance ‘dre42’ by Algorithm 1 exactly in 13 minutes.

Table 2: Numerical performance of the ‘dre’ problem instances [18]

Problem	opt	PDCA	gap (%)	time	permutation/bound
dre15	306	306	0	11	1 13 4 6 7 9 11 5 12 14 1 15 10 2 3 8
dre18	332	332	0	16	4 14 18 9 10 12 2 15 7 3 5 8 6 11 13 17 1 16
dre21	356	356	0	33	5 8 17 18 12 13 1 11 3 9 16 4 6 20 7 19 14 10 15 2 21
dre24	396	396	0	15	3 23 14 21 22 10 16 9 7 5 8 18 13 4 2 17
Continued on next page					

Table 2 – continued from previous page

Problem	opt	PDCA	gap(%)	time	permutation/bound
					1 19 12 11 15 24 6 20
dre30	508	508	0	1:22	28 2 1 17 6 3 11 21 19 22 24 8 26 20 23 13 4 29 18 25 10 30 16 15 14 12 7 5 27 9
dre42	764	764	0	13:00	3 36 41 28 30 14 34 32 42 37 33 10 27 12 35 9 7 21 5 29 18 11 8 38 24 2 15 22 6 1 13 19 40 23 25 39 31 16 17 26 4 20

In Table 3, the upper bounds generated by Algorithm 1 are compared with the state-of-art optimal values (or the best known upper bounds) in QAPLIB. Except *burxxx* and *skoxx* cases, we find that most instances can either be solved exactly or achieve an upper bound, which is accurate up to a relative error of 5% through the penalized DC relaxation. Because the subproblems of the corresponding penalized DC problems are failed to achieve the stopping criteria 10^{-6} of SDPNAL+, Algorithm 1 only provides the feasible solutions for *burxxx* cases. We note that the QAPLIB bounds were typically achieved using a rather large collection of different algorithms, which generally involve a branch and bound procedure requiring multiple convex relaxations, while our results are achieved by using a single relaxation.

Table 3: Numerical performance of the QAPLIB instances

Problem	opt	PDCA	gap (%)	time	permutation/bound
bur26a	5426670	5566175	2.57	2:33	11 26 15 7 4 13 12 6 2 18 1 9 5 21 8 14 3 19 20 17 10 25 24 16 23 22
bur26b	3817852	3956961	3.64	1:55	15 16 10 7 4 12 2 23 22 18 5 9 1 21 8 14 3 20 19 25 11 26 24 17 6 13
bur26c	5426795	5523812	1.79	7:03	13 3 12 7 16 11 25 10 15 9 8 19 18 20 4 21 1 14 5 6 22 24 2 23 26 17
bur26d	3821225	3902248	2.12	3:59	22 23 3 2 16 11 17 21 15 9 8 18 19 20 12 25 14 1 5 13 24 6 4 7 26 10
bur26e	5386879	5470899	1.56	6:26	22 3 6 7 12 26 1 16 11 9 18 19 20 14 13 8 5 15 21 2 17 24 4 10 25 23
bur26f	3782044	3847551	1.73	6:16	6 22 4 3 12 25 7 1 23 15 20 18 19 14 16 10 5 21 9 24 2 17 26 13 11 8
bur26g	10117172	10332466	2.13	7:32	2 11 22 23 13 10 25 8 1 21 20 4 7 18 12 15 9 19 5 26 16 6 14 3 24 17
bur26h	7098658	7257159	2.23	4:06	22 16 13 26 14 10 21 1 8 15 4 20 18 7 12 17 19 5 9 2 11 3 23 6 24 25
chr12a	9552	9552	0	04	7 5 12 2 1 3 9 11 10 6 8 4
chr12b	9742	9742	0	04	5 7 1 10 11 3 4 2 9 6 12 8
chr12c	11156	11156	0	04	7 5 1 3 10 4 8 6 9 11 2 12
chr15a	9896	9896	0	07	5 10 8 13 12 11 14 2 4 6 7 15 3 1 9
chr15b	7990	7990	0	07	4 13 15 1 9 2 5 12 6 14 7 3 10 11 8
chr15c	9504	9504	0	07	13 2 5 7 8 1 14 6 4 3 15 9 12 11 10

Continued on next page

Table 3 – continued from previous page

Problem	opt	PDCA	gap(%)	time	permutation/bound
chr18a	11098	11098	0	13	3 13 6 4 18 12 10 5 1 11 8 7 17 14 9 16 15 2
chr18b	1534	1534	0	1:52	10 7 11 8 12 9 15 6 14 5 13 4 16 1 17 2 18 3
chr20a	2192	2192	0	27	3 20 7 18 9 12 19 4 10 11 1 6 15 8 2 5 14 16 13 17
chr20b	2298	2298	0	28	20 3 9 7 1 12 16 6 8 14 10 4 5 13 17 2 18 11 19 15
chr20c	14142	14142	0	28	12 6 9 2 10 11 3 4 15 18 7 13 16 5 14 17 19 1 8 20
chr22a	6156	6156	0	39	15 2 21 8 16 1 7 18 14 13 5 17 6 11 3 4 20 19 9 22 10 12
chr22b	6194	6194	0	28	10 19 3 1 20 2 6 4 7 8 17 13 11 15 21 12 9 5 22 14 18 16
chr25a	3796	3796	0	2:00	25 12 5 3 18 4 16 8 20 10 14 6 23 15 24 19 13 1 21 11 17 2 22 7 9
els19	17212548	17212548	0	31	9 10 7 19 14 18 13 17 6 11 4 5 12 8 16 15 1 2 3
esc16a	68	68	0	18	12 15 7 11 14 6 10 8 4 16 13 3 5 9 2 1
esc16b	292	292	0	40	7 6 8 14 16 10 13 2 15 4 12 5 1 11 9 3
esc16c	160	160	0	1:00	15 10 9 2 1 14 11 3 4 13 8 16 12 6 7 5
esc16d	16	16	0	8	14 6 7 10 13 5 16 2 4 1 3 12 15 11 8 9
esc16e	28	30	7.14	15	15 10 7 14 4 8 6 16 12 2 1 5 9 13 11 3
esc16g	26	26	0	13	7 12 10 16 4 8 6 14 11 3 2 1 15 13 5 9
esc16h	996	996	0	18	6 5 13 7 12 11 15 4 8 3 16 9 2 1 10 14
esc16i	14	14	0	19	7 5 3 1 9 10 12 2 4 6 11 13 15 8 14 16
esc16j	8	8	0	11	11 4 5 8 14 16 13 9 7 1 10 12 15 3 6 2
esc32a	130	150	15.38	13:11	28 12 27 19 4 18 16 21 11 32 14 8 26 10 25 23 6 9 13 17 15 22 7 31 30 24 29 5 3 1 20 2
esc32b	168	168	0	12:18	1 3 9 25 11 27 5 7 6 2 8 4 13 29 14 10 30 26 15 31 16 12 32 28 21 19 23 20 18 17 22 24
esc32c	642	646	0.62	2:26:27	11 30 14 15 17 24 32 1 23 4 2 18 22 6 21 5 8 7 20 19 29 9 3 13 26 12 16 31 25 27 10 28
esc32e	2	2	0	12:36	19 2 17 9 3 7 12 30 16 10 14 13 31 15 21 18 22 28 4 26 6 25 24 11 20 8 27 5 1 32 23 29
esc32g	6	6	0	12:23	2 13 5 6 14 22 20 15 16 26 28 8 32 23 31 1 11 17 12 24 3 7 19 4 30 25 21 10 27 29 18 9
had12	1652	1652	0	4	3 10 11 2 12 5 7 6 8 1 4 9
had14	2724	2724	0	5	8 13 10 5 12 11 2 14 3 6 7 1 9 4
had16	3720	3720	0	8	9 4 16 1 7 8 6 14 15 11 12 10 5 3 2 13
had18	5358	5358	0	11	8 15 16 14 7 18 6 11 1 10 12 5 3 13 2 17 9 4
had20	6922	6922	0	42	8 15 1 14 19 6 7 17 16 12 10 11 5 20 2 3 4 9 18 13
kra30a	88900	88900	0	23:16	9 13 28 27 8 7 10 30 20 21 23 19 24 29 14 1 11 12 18 16 17 22 26 2 5 3 25 6 4 15
kra30b	91420	92010	0.65	14:45	24 15 22 6 4 3 5 2 17 1 16 11 25 26 30 19 10 28 29 21 23 20 8 12 9 7 13 27 18 14
kra32	88900	89100	0.22	2:11:24	8 10 5 9 6 14 27 23 24 31 12 16 28 4 13 7 11 3 29 22 26 15 30 21 1 2 20 32 25 18 19 17

Continued on next page

Table 3 – continued from previous page

Problem	opt	PDCA	gap(%)	time	permutation/bound
lipa20a	3683	3683	0	18	19 17 7 1 5 9 10 12 4 16 20 6 3 14 11 15 13 8 2 18
lipa20b	27076	27076	0	19	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20
lipa30a	13178	13178	0	4:43	9 13 22 17 25 23 29 12 11 6 5 28 20 27 14 4 18 8 19 30 21 7 15 24 26 3 16 1 10 2
lipa30b	151426	151426	0	4:51	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30
lipa40a	31538	31538	0	34:01	7 6 14 27 19 37 21 2 16 1 40 24 30 23 5 28 22 8 20 35 32 26 29 3 4 11 36 10 13 38 9 17 31 18 33 15 25 34 39 12
lipa40b	476581	476581	0	40:58	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40
lipa50a	62093	62093	0	20:30	28 32 37 39 49 23 19 44 33 7 14 30 15 5 36 6 17 26 48 25 40 3 45 27 18 31 29 16 9 12 1 8 4 2 50 21 43 35 24 38 34 46 42 13 20 22 41 47 10 11
lipa50b	1210244	1210244	0	23:04	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50
lipa60a	107218	107218	0	2:44:09	25 48 17 4 50 13 16 41 1 37 22 27 46 34 38 12 9 3 8 33 47 54 31 43 40 10 35 23 29 57 2 6 51 56 49 21 30 36 15 39 59 18 52 28 26 44 14 60 32 5 20 11 55 45 53 24 42 7 58 19
lipa60b	2520135	2520135	0	36:07	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60
nug12	578	578	0	12	3 9 7 12 1 11 8 4 2 10 6 5
nug14	1014	1014	0	4:40	9 8 13 2 1 11 7 14 3 4 12 5 6 10
nug15	1150	1150	0	5:19	12 5 6 15 10 11 7 14 3 4 9 8 13 2 1
nug16a	1610	1610	0	32	9 14 2 15 16 3 10 12 8 11 6 5 7 1 4 13
nug16b	1240	1240	0	9:27	8 11 3 5 13 9 7 1 12 2 10 6 16 4 15 14
nug17	1732	1732	0	10:17	16 15 2 14 9 11 8 12 10 3 4 1 7 6 13 17 5
nug18	1930	1936	0.31	29	17 1 7 5 6 11 8 12 10 13 4 15 2 3 9 16 18 14
nug20	2570	2570	0	2:40	18 14 10 3 9 4 2 12 11 16 19 15 20 8 13 17 5 7 1 6
nug21	2438	2442	0.16	17:25	4 18 11 16 13 6 15 14 19 8 7 1 12 17 20 21 3 9 10 2 5
nug22	3596	3642	1.28	4:02	20 4 16 8 19 7 10 9 13 21 17 14 15 11 3 18 22 1 12 6 2 5
nug25	3744	3750	0.16	35:06	12 4 14 23 13 24 21 7 10 20 17 16 6 19 1 11 25 8 9 15 18 2 3 22 5
nug27	5234	5236	0.04	30:21	23 18 3 1 27 17 5 13 7 15 12 26 8 19 20 2 24 21 4 10 9 14 22 25 6 16 11
nug28	5166	5166	0	35:33	11 8 20 28 1 9 18 26 16 17 19 10 15 7 14 27

Continued on next page

Table 3 – continued from previous page

Problem	opt	PDCA	gap(%)	time	permutation/bound
					4 13 25 6 22 12 5 3 24 2 21 23
rou12	235528	235528	0	04	6 5 11 9 2 8 3 1 12 7 4 10
rou15	354210	354210	0	5:46	12 6 8 13 5 3 15 2 7 1 9 10 4 14 11
rou20	725522	725582	0	4:54	12 11 9 17 7 14 8 13 6 10 19 18 5 2 16 3 4 1 20 15
scr12	31410	31410	0	04	8 6 11 10 2 9 5 1 12 7 4 3
scr15	51140	51140	0	06	12 10 11 14 1 13 9 5 15 6 4 2 8 7 3
scr20	110030	110030	0	13:42	20 7 12 6 4 8 3 2 14 11 18 9 19 15 16 17 13 5 10 1
sko42	15812	15952	0.89	3:58:53	23 16 30 36 3 21 8 22 39 37 24 1 38 6 28 2 14 18 32 40 25 33 4 35 26 9 15 29 27 10 19 20 34 42 11 13 31 5 12 7 17 41
sko56	34458	36490	5.90	3:18:03	21 42 2 11 30 51 49 3 25 6 56 33 9 23 13 54 38 24 8 50 32 36 44 10 31 45 53 37 27 12 14 41 43 55 22 5 29 20 52 35 26 46 4 28 34 48 18 40 19 15 16 1 47 17 7 39
sko64	48498	50948	5.05	3:48:56	31 57 56 53 51 3 43 15 2 54 17 52 23 32 11 27 63 26 59 49 13 6 40 46 33 7 44 16 64 39 38 61 10 42 45 24 4 62 30 41 35 37 1 21 55 5 28 34 9 25 58 22 19 8 48 12 18 50 14 20 29 47 60 36
sko72	66256	70318	6.13	14:42:49	45 11 51 71 29 14 60 31 12 53 41 50 57 44 28 21 10 64 25 52 68 1 6 13 47 67 35 43 20 16 59 49 7 22 23 27 32 4 63 55 33 46 65 17 30 24 69 66 36 9 38 37 2 34 3 72 42 19 39 61 26 62 56 54 5 58 15 40 48 70 8 18
sko81	90998	93356	2.59	25:47:17	47 67 80 65 34 39 69 74 30 40 23 63 38 20 33 26 81 25 54 4 79 35 51 2 12 14 3 77 24 73 58 32 70 21 78 61 10 19 75 37 18 15 49 42 72 22 43 59 13 44 7 41 6 28 71 55 62 1 11 31 5 36 9 53 29 46 52 27 45 56 17 50 8 60 64 76 16 68 57 48 66
ste36a	9526	9640	1.20	1:12:45	35 16 1 15 14 28 29 30 31 17 18 10 7 11 20 19 32 34 2 8 4 13 12 23 22 21 33 36 3 9 5 6 27 26 25 24
ste36b	15852	15932	0.50	25:14	35 31 30 29 28 1 15 9 16 33 34 32 19 20 7 10 18 17 26 25 23 14 11 13 4 8 2 24 22 21 27 12 6 5 3 36
ste36c	8239110	8254628	0.19	2:19:21	24 25 26 27 11 6 5 3 35 22 21 23 14 12 13 4 8 2 33 32 19 28 20 7 10 18 17 34 31 30 29 15 1 9 16 36
tai12a	224416	224416	0	05	8 1 6 2 11 10 3 5 9 7 12 4
tai12b	39464925	39464925	0	4	9 4 6 3 11 7 12 2 8 10 1 5
tai15a	388214	388870	0.17	31	6 10 4 7 2 9 1 11 3 14 13 15 5 12 8
tai15b	51765268	51765268	0	15	1 9 4 6 8 15 7 11 3 5 2 14 13 12 10
tai17a	491812	491812	0	6:18	12 2 6 7 4 8 14 5 11 3 16 13 17 9 1 10 15
tai20a	703482	703482	0	14:48	10 9 12 20 19 3 14 6 17 11 5 7 15 16 18 2 4 8 13 1
tai20b	122455319	122455319	0	48	8 16 14 17 4 11 3 19 7 9 1 15 6 13 10 2

Continued on next page

Table 3 – continued from previous page

Problem	opt	PDCA	gap(%)	time	permutation/bound
					5 20 18 12
tai25a	1167256	1217842	4.33	31:47	20 1 10 7 9 13 4 19 3 2 11 15 8 21 12 14 18 25 23 17 5 22 24 6 16
tai25b	344355646	344855160	0.15	16:03	4 25 16 9 13 5 6 19 7 17 10 3 15 20 18 2 22 23 8 11 21 24 14 12 1
tai30a	1818146	1818146	0	33:33	19 18 4 24 30 25 5 7 1 22 28 20 11 13 9 16 8 10 17 21 12 29 2 15 3 14 26 27 23 6
tai30b	637117113	644555585	1.17	4:52	4 15 5 8 21 11 30 14 17 20 6 13 18 7 23 10 24 27 29 9 19 28 2 26 12 22 25 16 1 3
tai35a	2422002	2431214	0.38	37:01	35 9 21 7 23 2 12 8 20 26 33 16 18 24 22 15 30 3 14 11 1 5 17 4 10 13 27 29 34 32 25 31 28 6 19
tai35b	283315445	284614706	0.46	1:14:45	1 17 5 30 11 2 22 3 14 10 9 20 18 12 19 33 8 32 13 27 15 21 34 35 7 24 6 23 31 28 16 4 29 26 25
tai40a	3139370	3154106	0.47	1:54:41	6 14 15 12 9 3 19 20 28 27 5 2 7 31 36 4 37 13 29 35 38 32 11 1 22 39 18 30 23 33 25 24 21 8 10 17 34 16 40 26
tai40b	637250948	640933239	0.46	2:01:44	19 1 25 11 37 31 36 15 39 13 22 7 40 27 4 33 14 34 9 10 38 23 5 32 35 12 16 3 24 2 21 28 20 17 26 6 30 29 18 8
tai50a	4938796	5086610	2.99	11:54:27	5 42 35 34 36 37 4 7 21 10 8 25 48 27 41 11 18 32 6 44 12 33 15 2 29 50 20 3 43 14 46 38 26 40 47 39 19 28 45 49 13 30 23 1 16 22 17 31 9 24
tai60b	7205962	7417848	2.94	5:41:01	18 41 38 7 20 26 30 36 16 31 59 57 21 54 19 48 23 47 27 4 10 14 5 43 58 3 6 50 12 49 8 44 17 29 15 22 33 46 35 32 25 52 34 39 45 55 60 11 1 13 53 37 51 9 40 28 24 2 42 56
tho30	149936	151156	0.81	26:26	29 14 1 2 8 28 22 25 12 20 19 11 24 27 17 26 30 10 6 15 3 7 5 4 16 23 21 13 9 18
tho40	240516	242282	0.73	5:38:41	38 37 5 13 26 27 35 31 4 28 9 32 21 8 29 25 18 33 22 16 30 6 12 34 39 14 20 15 1 10 11 17 19 2 40 23 24 7 36 3

6 Conclusion

This paper established an exact rank constrained DNN formulation of QAP. Under the framework of DC programming, we can efficiently solve the penalized DC problem using the semi-proximal augmented Lagrangian method. If the subproblems can be solved successfully, our algorithm usually reaches the optimal solutions of QAP exactly. Even if the subproblem is challenging to solve, our proposed algorithm can still provide a good feasible solution close to the optimal upper bound in QAPLIB. As future work, we will investigate the structure of the constraints of the penalized DC problem and try to reduce

the number of constraints for solving the rank constrained DNN formulation of QAPs more efficiently.

Acknowledgements We would like to thank Dr. Xudong Li and Dr. Ying Cui for many helpful discussions on this work. Also, we would like to thank the two anonymous referees and the associate editor for their valuable comments and constructive suggestions which have helped to improve the quality of this paper.

References

1. An, L.T.H., Tao, P.D.: DC programming and DCA: thirty years of developments, *Mathematical Programming* **169**, 5-68 (2018)
2. An, L.T.H., Tao, P.D., Huynh, V.N.: Exact penalty and error bounds in DC programming, *Journal of Global Optimization* **52**, 509-535 (2012)
3. Anstreicher, K.: Recent advances in the solution of quadratic assignment problems, *Mathematical Programming* **97**, 27-42 (2003)
4. Anstreicher, K., Wolkowicz, H.: On Lagrangian relaxation of quadratic matrix constraints, *SIAM Journal on Matrix Analysis and Applications* **22**, 41-55 (2000)
5. Attouch, H., Bolte, J.: On the convergence of the proximal algorithm for nonsmooth functions involving analytic features, *Mathematical Programming*, **116**, 5-16 (2009).
6. Attouch, H., Bolte, J., Redont, P., Soubeyran, A.: Proximal Alternating Minimization and Projection Methods for Nonconvex Problems: An Approach Based on the Kurdyka-Lojasiewicz Inequality, *Mathematics of Operations Research* **35**, 438-457 (2010)
7. Ben-Tal, A., Nemirovski, A.: *Lectures on Modern Convex Optimization: analysis, algorithms, and engineering applications*, vol. 2, Society for Industrial Mathematics, 2001.
8. Bi, S.J., Pan, S.H.: Error bounds for rank constrained optimization problems and applications, *Operations Research Letters* **44**, 336-341 (2016)
9. Bolte, J., Daniilidis, A., Lewis, A.S.: The Lojasiewicz inequality for nonsmooth subanalytic functions with applications to subgradient dynamical systems. *SIAM Journal on Optimization* **17**, 1205-1223 (2007)
10. Bolte, J., Daniilidis, A., Lewis, A.S., Shiota, M.: Clarke subgradients of stratifiable functions. *SIAM Journal on Optimization* **18**, 556-572 (2007).
11. Bolte, J., Pauwels, E.: Majorization-minimization procedures and convergence of SQP methods for semi-algebraic and tame programs, *Mathematics of Operations Research*, **41**, 442-465 (2016).
12. Bolte, J., Sabach, S., Teboulle, M.: Proximal alternating linearized minimization for nonconvex and nonsmooth problems, *Mathematical Programming*, **146**, 459-494 (2014).
13. Burer, S.: On the copositive representation of binary and continuous nonconvex quadratic programs, *Mathematical Programming* **120**, 479-495 (2009)
14. Burkard, P.: Quadratic assignment problems, in *Handbook of Combinatorial Optimization*, Pardalos, P.M., Du, D.Z., Graham, R.L. (ed.), 2741-2814, Springer, New York (2013)
15. Buss, F., Frandsen, G. S., Shallit, J.O.: The computational complexity of some problems of linear algebra, *Journal of Computer and System Sciences* **58**, 572-596 (1999)
16. Dontchev, A.L., Rockafellar, R.T.: *Implicit functions and solution mappings - a view from variational analysis*, Springer, (2009)
17. Drezner, Z.: *The quadratic assignment problem*, Location Science, 345-363, Springer, New York (2015)
18. Drezner, Z., Hahn, P., Taillard, É.D.: Recent advances for the quadratic assignment problem with special emphasis on instances that are difficult for meta-heuristic methods, *Operation Research* **139**, 65-94 (2005)
19. Coste, M.: *An Introduction to o-minimal Geometry*, RAAG Notes, Institut de Recherche Mathématiques de Rennes, 1999.
20. Fu, T., Ge, D., Ye, Y.: On doubly positive semidefinite programming relaxations, *Journal of Computational Mathematics* **36**, 391-403 (2018)
21. Gao, Y.: *Structured Low Rank Matrix Optimization Problems: A Penalized Approach*, PhD thesis, National University of Singapore (2010)

22. Gao, Y., Sun, D.F.: A majorized penalty approach for calibrating rank constrained correlation matrix problems, Preprint available at http://www.mypolyuweb.hk/~dfsun/MajorPen_May5.pdf (2010)
23. Hahn, P., Anjos, M.: QAPLIB - a quadratic assignment problem library, <http://www.seas.upenn.edu/qaplib>.
24. Horn, R.A., Johnson, C.R.: Matrix Analysis, Cambridge University Press, New York (1985)
25. Ioffe, A. D.: An Invitation to tame optimization, *SIAM Journal on Optimization* **19**, 1894-1917 (2009)
26. Kim, S., Kojima, M., Toh, K.C.: A Lagrangian-DNN relaxation: a fast method for computing tight lower bounds for a class of quadratic optimization problems, *Mathematical Programming* **156**, 161-187 (2016)
27. Koopmans, T.C., Beckmann, M.J.: Assignment problems and the location of economics activities, *Econometrica* **25**, 53-76 (1957)
28. Li, Q., Qi, H.-D.: A Sequential Semismooth Newton Method for the Nearest Low-rank Correlation Matrix Problem. *SIAM Journal on Optimization* **21**, 1641-1666 (2011).
29. Lin, C.-J., Saigal, R.: On solving large-scale semidefinite programming problems a case study of quadratic assignment problem. Technical report, Department of Industrial and Operations Engineering, University of Michigan, Ann Arbor MI, (1997)
30. Liu, T., Pong, T.K., Takeda, A.: A refined convergence analysis of with applications to simultaneous sparse recovery and outlier detection. *Computational Optimization Applications* **73**, 69-100 (2019).
31. Mishra, B.: Algorithmic algebra. Springer-Verlag New York, 1993.
32. Motzkin, T.S., Straus, E.G.: Maxima for graphs and a new proof of a theorem of Turan, *Canadian Journal of Mathematics* **17**, 533-540 (1965)
33. Murty, K.G., Kabadi, S.N.: Some NP-complete problems in quadratic and nonlinear programming, *Mathematical Programming* **39**, 117-129 (1987)
34. Povh, J., Rendl, F.: A copositive programming approach to graph partitioning, *SIAM Journal on Optimization* **18**, 223-241 (2007)
35. Povh, J., Rendl, F.: Copositive and semidefinite relaxations of the quadratic assignment problem, *Discrete Optimization* **6**, 231-241 (2009)
36. Ramana, M., Tunçel, L., Wolkowicz, H.: Strong duality for semidefinite programming, *SIAM Journal on Optimization* **7**, 641-662 (1997)
37. Rendl, F., Sotirov, R.: Bounds for the quadratic assignment problem using the bundle method, *Mathematical Programming* **109**, 505-524 (2007)
38. Rockafellar, R.T.: *Convex Analysis*, Princeton University Press, Princeton (1970)
39. Sahni, S., Gonzalez, T.: P-complete approximation problems, *Journal of the ACM* **23**, 555-565 (1976)
40. Sun, D.F., Toh, K.C., Yuan, Y.C., Zhao, X.Y.: SDPNAL+: A Matlab software for semidefinite programming with bound constraints (version 1.0), *Optimization Methods and Software*, in print (2019)
41. Pham, D.T., Le Thi, H.A.: Convex analysis approach to DC programming: theory, algorithms and applications. *ACTA Mathematica Vietnamica* **22**, 289-355 (1997)
42. Pham, D.T., Le Thi, H.A.: ADC optimization algorithm for solving the trust-region subproblem. *SIAM Journal on Optimization* **8**, 476-505 (1998)
43. Le Thi, H.A., Pham, D.T.: DC programming and DCA: thirty years of developments. *Mathematical Programming* **169**, 5-68 (2018)
44. Todd, M.J.: Semidefinite optimization. *Acta Numerica*. **10**, 515-560 (2001).
45. Vandenberghe, L., Boyd, S.: Semidefinite programming, *SIAM Review* **38**, 49-75 (1996)
46. Wen, Z.W., Goldfarb, D., Yin, W.T.: Alternating direction augmented Lagrangian methods for semidefinite programming, *Mathematical Programming Computation* **2**, 203-230 (2010)
47. Weyl, H.: Das asymptotische verteilungsgesetz der eigenwerte linearer partieller differentialgleichungen (mit einer anwendung auf die theorie der hohlraumstrahlung, *Mathematische Annalen* **71**, 441-479 (1912)
48. Yang, L.Q., Sun, D.F., Toh, K.C.: SDPNAL+: A majorized semismooth Newton-CG augmented lagrangian method for semidefinite programming with nonnegative constraints, *Mathematical Programming Computation* **7**, 331-366 (2015)

-
49. Yoshise, A., Matsukawa, Y.: On optimization over the doubly nonnegative cone, Proceedings of 2010 IEEE Multi-conference on Systems and Control, 13-19 (2010)
 50. Zhao, Q., Karisch, S.E., Rendl, F., Wolkowicz, H.: Semidefinite programming relaxations for the quadratic assignment problem, Journal of Combinatorial Optimization **2**, 71-109 (1998)
 51. Zhao, X.Y., Sun, D.F., Toh, K.C.: A Newton-CG augmented lagrangian method for semidefinite programming, SIAM Journal on Optimization **20**, 1737-1765 (2010)